


Application No. 09/618,066
 Attorney Docket No. 5725.0656-00

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions and listings of claims in the application:

1-156. (Canceled)

157. (Currently amended) A process for ~~limiting the migration~~ non-migrating deposit of a lipstick composition comprising including in said lipstick composition at least one liquid continuous fatty phase, said at least one liquid continuous fatty phase being structured with a sufficient amount of an agent for ~~limiting the migration~~ non-migrating deposit of said lipstick composition, said agent comprising at least one structuring polymer, wherein said at least one structuring polymer is chosen from polymers of formula (I) below and mixtures thereof:



in which:

- n is an integer which represents the number of amide units such that the number of ester groups present in said at least one structuring polymer ranges from 10% to 50% of the total number of all said ester groups and all said amide groups comprised in said at least one structuring polymer;

- R¹, which are identical or different, are each chosen from alkyl groups comprising at least 4 carbon atoms and alkenyl groups comprising at least 4 carbon atoms;

Application No. 09/618,066
Attorney Docket No. 5725.0656-00

- R^2 , which are identical or different, are each chosen from C_4 to C_{42}

hydrocarbon-based groups with the proviso that at least 50% of R^2 are chosen from C_{30} to C_{42} hydrocarbon-based groups;

- R^3 , which are identical or different, are each chosen from organic groups comprising atoms chosen from carbon atoms, hydrogen atoms, oxygen atoms and nitrogen atoms with the proviso that R^3 comprises at least 2 carbon atoms; and

- R^4 , which are identical or different, are each chosen from hydrogen atoms, C_1 to C_{10} alkyl groups and a direct bond to group chosen from R^3 and another R^4 such that when said at least one group is chosen from another R^4 , the nitrogen atom to which both R^3 and R^4 are bonded forms part of a heterocyclic structure defined in part by R^4 -N- R^3 , with the proviso that at least 50% of all R^4 are chosen from hydrogen atoms; and

wherein said lipstick composition further comprises at least one dyestuff chosen from pigments and nacles.

158. (Previously presented) A process according to Claim 157, wherein said lipstick composition has a hardness ranging from 20 g to 2000 g.


159. (Original) A process according to Claim 158, wherein said hardness ranges from 20 g to 900 g.

160. (Original) A process according to Claim 159, wherein said hardness ranges from 20 g to 600 g.

161. (Currently amended) A process for ~~limiting the migration~~ non-migrating deposit of a lipstick composition comprising at least one continuous liquid fatty phase comprising structuring said fatty phase with a sufficient amount of at least one

Application No. 09/618,066
 Attorney Docket No. 5725.0656-00

structuring polymer, wherein said at least one structuring polymer is chosen from polymers of formula (I) below and mixtures thereof:



in which:

- n is an integer which represents the number of amide units such that the number of ester groups present in said at least one structuring polymer ranges from 10% to 50% of the total number of all said ester groups and all said amide groups comprised in said at least one structuring polymer;

- R^1 , which are identical or different, are each chosen from alkyl groups comprising at least 4 carbon atoms and alkenyl groups comprising at least 4 carbon atoms;

- R^2 , which are identical or different, are each chosen from C_4 to C_{42} hydrocarbon-based groups with the proviso that at least 50% of R^2 are chosen from C_{30} to C_{42} hydrocarbon-based groups;

- R^3 , which are identical or different, are each chosen from organic groups comprising atoms chosen from carbon atoms, hydrogen atoms, oxygen atoms and nitrogen atoms with the proviso that R^3 comprises at least 2 carbon atoms; and


- R^4 , which are identical or different, are each chosen from hydrogen atoms, C_1 to C_{10} alkyl groups and a direct bond to group chosen from R^3 and another R^4 such that when said at least one group is chosen from another R^4 , the nitrogen atom to which

Application No. 09/618,066
Attorney Docket No. 5725.0656-00

both R^3 and R^4 are bonded forms part of a heterocyclic structure defined in part by R^4 - N - R^3 , with the proviso that at least 50% of all R^4 are chosen from hydrogen atoms; and
wherein said lipstick composition further comprises at least one dyestuff chosen from pigments and nacres.

162-167. (Canceled)

168. (Currently amended) A process for ~~limiting the migration~~ non-migrating deposit of a foundation composition comprising including in said foundation composition at least one liquid continuous fatty phase, said at least one liquid continuous fatty phase being structured with a sufficient amount of an agent for ~~limiting the migration~~ non-migrating deposit of said foundation composition, said agent comprising at least one structuring polymer, wherein said at least one structuring polymer is chosen from polymers of formula (I) below and mixtures thereof:



in which:

- n is an integer which represents the number of amide units such that the number of ester groups present in said at least one structuring polymer ranges from 10% to 50% of the total number of all said ester groups and all said amide groups comprised in said at least one structuring polymer;

Application No. 09/618,066
Attorney Docket No. 5725.0656-00

- R^1 , which are identical or different, are each chosen from alkyl groups comprising at least 4 carbon atoms and alkenyl groups comprising at least 4 carbon atoms;

- R^2 , which are identical or different, are each chosen from C_4 to C_{42} hydrocarbon-based groups with the proviso that at least 50% of R^2 are chosen from C_{30} to C_{42} hydrocarbon-based groups;

- R^3 , which are identical or different, are each chosen from organic groups comprising atoms chosen from carbon atoms, hydrogen atoms, oxygen atoms and nitrogen atoms with the proviso that R^3 comprises at least 2 carbon atoms; and

- R^4 , which are identical or different, are each chosen from hydrogen atoms, C_1 to C_{10} alkyl groups and a direct bond to group chosen from R^3 and another R^4 such that when said at least one group is chosen from another R^4 , the nitrogen atom to which both R^3 and R^4 are bonded forms part of a heterocyclic structure defined in part by R^4 -N- R^3 , with the proviso that at least 50% of all R^4 are chosen from hydrogen atoms; and

wherein said foundation composition further comprises at least one dyestuff chosen from pigments and naces.


169. (Previously presented) A process according to Claim 168, wherein said foundation composition has a hardness ranging from 20 g to 2000 g.

170. (Previously presented) A process according to Claim 169, wherein said hardness ranges from 20 g to 900 g.

171. (Previously presented) A process according to Claim 170, wherein said hardness ranges from 20 g to 600 g.

Application No. 09/618,066
Attorney Docket No. 5725.0656-00

172. (Previously presented) A process for ~~limiting the migration~~ non-migrating deposit of a foundation composition comprising at least one continuous liquid fatty phase comprising structuring said fatty phase with a sufficient amount of at least one structuring polymer, wherein said at least one structuring polymer is chosen from polymers of formula (I) below and mixtures thereof:



in which:

- n is an integer which represents the number of amide units such that the number of ester groups present in said at least one structuring polymer ranges from 10% to 50% of the total number of all said ester groups and all said amide groups comprised in said at least one structuring polymer;

- R¹, which are identical or different, are each chosen from alkyl groups comprising at least 4 carbon atoms and alkenyl groups comprising at least 4 carbon atoms;


- R², which are identical or different, are each chosen from C₄ to C₄₂ hydrocarbon-based groups with the proviso that at least 50% of R² are chosen from C₃₀ to C₄₂ hydrocarbon-based groups;

- R³, which are identical or different, are each chosen from organic groups comprising atoms chosen from carbon atoms, hydrogen atoms, oxygen atoms and nitrogen atoms with the proviso that R³ comprises at least 2 carbon atoms; and

Application No. 09/618,066
Attorney Docket No. 5725.0656-00

- R^4 , which are identical or different, are each chosen from hydrogen atoms, C_1 to C_{10} alkyl groups and a direct bond to group chosen from R^3 and another R^4 such that when said at least one group is chosen from another R^4 , the nitrogen atom to which both R^3 and R^4 are bonded forms part of a heterocyclic structure defined in part by R^4 -N- R^3 , with the proviso that at least 50% of all R^4 are chosen from hydrogen atoms; and wherein said foundation composition further comprises at least one dyestuff chosen from pigments and nacres.

173. (New) A process for non-migrating deposit of a composition for making up at least one keratinous material comprising at least one continuous liquid fatty phase comprising structuring said fatty phase with a sufficient amount of at least one structuring polymer, wherein said at least one structuring polymer is chosen from polymers of formula (I) below and mixtures thereof:



in which:

- n is an integer which represents the number of amide units such that the number of ester groups present in said at least one structuring polymer ranges from 10% to 50% of the total number of all said ester groups and all said amide groups comprised in said at least one structuring polymer;

Application No. 09/618,066
Attorney Docket No. 5725.0656-00

- R^1 , which are identical or different, are each chosen from alkyl groups comprising at least 4 carbon atoms and alkenyl groups comprising at least 4 carbon atoms;

- R^2 , which are identical or different, are each chosen from C_4 to C_{42} hydrocarbon-based groups with the proviso that at least 50% of R^2 are chosen from C_{30} to C_{42} hydrocarbon-based groups;

- R^3 , which are identical or different, are each chosen from organic groups comprising atoms chosen from carbon atoms, hydrogen atoms, oxygen atoms and nitrogen atoms with the proviso that R^3 comprises at least 2 carbon atoms; and

- R^4 , which are identical or different, are each chosen from hydrogen atoms, C_1 to C_{10} alkyl groups and a direct bond to group chosen from R^3 and another R^4 such that when said at least one group is chosen from another R^4 , the nitrogen atom to which both R^3 and R^4 are bonded forms part of a heterocyclic structure defined in part by R^4 -N- R^3 , with the proviso that at least 50% of all R^4 are chosen from hydrogen atoms; and

wherein said composition for making up at least one keratinous material further comprises at least one dyestuff chosen from pigments and nacles.

174. (New) A process according to claim 173, wherein said at least one structuring polymer is chosen from ethylenediamine/stearyl dimer tallate copolymer.